

GEOHELMINTHS EGGS CONTAMINATION OF SANDPITS IN VLADIVOSTOK, RUSSIA.


Moskvina Tatyana Vladimirovna*

Bartkova Albina Dmitrievna¹


Ermolenko Alexey Vasil'evich²

The most widespread geohelminths

? *Toxocara canis* life cycle



? *Ascaris lumbricoides* life cycle



Materials and methods

- ? Totally 60 samples from five district were collected and analyzed. Sand samples were analyzed using Romanenko's method




NaNO_3

Results




Egg with larva


Toxocara spp.eggs



Ascaris sp.egg



Toxascaris leonina egg



Results

Table 1 Parasites found in soil samples

Parasites found	Number of soil samples	Min-max number of eggs/25 gr
<i>Toxascaris leonina</i>	2	1-23
<i>Toxocara</i>	14	1-6
<i>Ascaris</i> sp.	7	1-3

Table 2 Results of sandpits samples analyses in each district

District	Total number of collected samples	Number of positive samples	Min-max number of eggs/25 g
Pervorechenskij	12	3	3-5
Sovetskij	11	3	1-3
Pervomajskij	16	5	1-26
Frunzenskij	11	4	1-3
Leninskii'	10	3	2-18



Control and prevention



Thanks' for your attention