

The Unique Australian Wildlife

**The notes to the geological history of the
continent**

By Maria Skochilova

School 7

Nizhny Tagil



Many countries have their own
unique fauna.

But Australia is mostly unusual
in that

320-270 million years ago

Permo-Carboniferous Age



If we had observed the Earth surface from space at that time, we would have seen quite the other picture

270-210 million years ago


The end of Permian – the beginning of Trias
About 275 million years ago

Euroamerica and
Angara
made a huge landmass

The Tethys Sea
Separated Laurasia

from the
protocontinent

of Pangaea.
Nowadays it is
the Middeteranian
Sea




270-210 million years ago

The end of Permian – the beginning of Trias

- ❖ On land the vertebrates are represented in the Triassic by amphibians and reptiles.
- ❖ The first true mammals, which were very small, are supposed to appear in the Late Triassic.

Triassic Period:

continents and oceans of the Earth in Early Triassic time



Monotreme



**short-beaked
echidna**

**amphibious
platypus**


The egg-laying
mammalians include
the amphibious
platypus
and the terrestrial
echidnas
of Australia, Tasmania,
and New Guinea



180 million years ago

Middle Jurassic period



The protocontinent supposedly covered about half the Earth and was completely surrounded by a world ocean called Panthalassa.




180 million years ago

Middle Jurassic period

- ❖ Dinosaurs and other reptiles emerged to dominate the land, sea, and sky.
- ❖ The first birds and new varieties of reefbuilding and other invertebrate faunas, provided Jurassic life with added complexity.




Late Jurassic Epoch: geochronological map



100 million years ago

Early Cretaceous Period



Later Pangaea began to break apart. Its segments Laurasia and Gondwanaland gradually receded, resulting in the formation of the Atlantic Ocean.

100 million years ago

Early Cretaceous Period

Two important groups of modern mammals evolved during the Cretaceous.

Cretaceous placentals, smaller than those of present-day ones, were poised to take over the terrestrial environments as soon as the dinosaurs vanished.

100 million years ago


Early Cretaceous Period

Another mammal group, the marsupials, evolved during the Cretaceous as well.

This group includes the native species of Australia, kangaroos, koalas, and the North American opossum.



Late Cretaceous Epoch: geochronological map




70 million years ago

The end of Cretaceous Period



The Late Cretaceous record is much more complete. It is known, for instance, that during the Late Cretaceous many dinosaur types lived in relationships like the present-day terrestrial mammals.


Diprotodon



extinct marsupial mammals
existed 30 - 10,000 years
ago
in Australia.

- ❖ characterized by a wombat-like body the size of a large rhinoceros.
- ❖ massively constructed skeleton to support its imposing bulk.
- ❖ well developed teeth of gnawing animals.
- ❖ herbivorous
- ❖ distantly related to kangaroos and wombats.

45 million years ago By that time
The beginning of Cenozoic era Australasia was
isolated
from all other
continental masses,
here marsupials
evolved
into many diverse
forms.




In South America
they survived
alongside placentals,
forming the
Neotropical





Structural and behavioral parallels with placental mammals are in some cases quite striking.

Parallel evolution of marsupial and placental mammals

▲ marsupial
● placental



There are marsupials that look remarkably like moles, shrews, squirrels, mice, dogs, and hyenas.



The koala and the kangaroo
are the most well-known
marsupials.



Marsupials

Long-nosed
bandicoot



Spotted-tailed
quoll,
or native cat



Marsupials

Virginia,
or opossum



Marsupials



Red kangaroo

—
Wallaby



Western grey kangaroo

Marsupials



Kangaroo Rat



Dunnart,
a marsupial
mouse

Marsupials



Tasmanian Devil





Wombat

Marsupials
The niches that marsupials
fill
are closely associated
with structure.




The diets of marsupials
are as varied
as the niches they occupy.

The burrowing species have powerful foreclaws with which they can tunnel into the ground for food and for shelter



The gliders have a membrane along either flank, attached to the forelegs and hind legs, that enables the animals to glide down from a high perch

Cenozoic Era: faunal migration routes and barriers



The earliest isolation of Australia from all the other continents made its fauna unique



Literature

- Т. Клементьева, Дж. Шэннон Гарри English-3 – Обнинск: Титул, 2005
- Д. Эттенборо Живая природа – М.: Мир книги, 2001
- Britannica 2007 Ultimate Reference Suite DVD - энциклопедия, англоязычное издание: www.britannica.co.uk