

ARCH and GARCH

Modeling Volatility Dynamics

Modeling Unequal Variability

- **Equal Variability: Homoscedasticity**
- **Unequal Variability: Heteroscedasticity**
 - **Means any variability (around the mean) that is not homoscedasticity**
 - **Models must be developed for specific cases**

What These Acronym Mean?

- **ARCH**
 - Autoregressive Conditional Heteroscedasticity
- **GARCH**
 - Generalized ARCH

Information in e^2

- Let ε_t have the mean 0 and the variance σ_t .
- Let e_t be the residual of a model fitted.
- Then:
 - e_t estimates ε_t
 - e_t^2 estimates the variance σ_t^2 .

ARCH Modeling of σ_t^2 .

- **ARCH(1)**

$$\sigma_t^2 = \omega + \alpha \varepsilon_{(t-1)}^2$$

- **ARCH as AR(1) on** $\varepsilon_t^2 = \sigma_t^2 + v_t$

$$\varepsilon_t^2 = \omega + \alpha \varepsilon_{(t-1)}^2 + v_t$$

GARCH

- **GARCH(1)**

$$\sigma_t^2 = \omega + \alpha \varepsilon_{(t-1)}^2 + \beta \sigma_{(t-1)}^2$$

- **GARCH (1) as ARMA(1,1) on** $\varepsilon_t^2 = \sigma_t^2 + v_t$

$$\varepsilon_t^2 = \omega + (\alpha + \beta) \varepsilon_{(t-1)}^2 + v_t - \beta v_{(t-1)}$$

Asymmetry in GARCH - TARCH

- TARCH(1,1)

$$\sigma_t^2 = \omega + \alpha \varepsilon_{t-1}^2 + \gamma d \varepsilon_{t-1s}^2 + \beta \sigma_{t-1}^2$$

d = 1 if $\varepsilon_t < 0$, and = 0 if $\varepsilon_t \geq 0$

Asymmetry in GARCH - EGARCH

- **EGARCH(1,1)**

$$\log(\sigma_t^2) = \omega + \beta \log \sigma_{t-1}^2 + \alpha \left| \frac{\varepsilon_{t-1}}{\sigma_{t-1}} \right| + \gamma \frac{\varepsilon_{t-1}}{\sigma_{t-1}}$$

$\sigma_t^2 > 0$

$\gamma \neq 0$ for asymmetric effect

Eviews Command

ARCH(p, q) *series_name* c