

Inflammatory reaction of an organism


is carried out by a loose collagen tissue together with leukocytes of blood

- At damage of a tissue **leucocytes** react the first and throw out the granules in tissue, histamin increases permeability of capillaries for blood plasma and leukocytes. Plasma collects in a tissue and creates **an inflammatory oedema** which separates the inflammatory focus from healthy tissues and does not allow to extend to products of disintegration on an organism.
- **Neutrophils** come the first to the center of damage. They throw out in a tissue much oxygen superoxide, lysosomal enzymes and contents of specific granules, and then perish. Oxygen superoxide and lysocim destroy bacteria, cation-proteins increase an oedema, and lysosomal enzymes even more destroy the damaged tissue. This phase of an inflammation refers to **leucocytic** (or the **sharp period**).
- Then **the macrophagic phase** develops. **Macrophages** come to the center of damage and phagocyte bacteria and products of disintegration of tissue, completely clearing from them the inflammatory focus.
- **The fibroblastic (reparative) phase** begins. **Fibroblasts** are actively made multiple copies, form around of the cleared site of damage a fibroblastic capsule and secrete of fundamantal substance and collagen of 1 type of which at once form thick fibres, filling with them defect of a tissue. Vessels grow into a capsule, the granulation tissue is formed of thick, rough collagenic fibres and vessels. Within the next 6 months rough scar is gradually reconstructed, thick fibres from collagen of 1 type are replaced with more thin fibres from collagen 3 types

Лейкоцитарная фаза воспаления (в коже) Leucocytic phase of inflammation (in skin)



Макрофагическая фаза *Macrophagic phase*



Phagocytosis
of bacteria and
damaged tissue

Фагоцитоз
продуктов
распада



Destruction of AG,
Stimulation of restoration



Уничтожение АГ,
стимуляция процессов
восстановления

Cellular infiltration

Клеточный
инfiltrат

Фибробластическая (репаративная) фаза

Fibroblastic (reparative) phase

Поверхность раны

Surface of wound

Фибробластическая
капсула

Fibroblastic capsule



1. Образование капсулы из фибробластов.
Formation of capsule from fibroblasts
2. Секреция основного вещества и волокон.
Secretion of fundamental substance and fibers

Формирование рубца

Formation of scar



Рубец

Scar

Грануляционная
ткань

Granulation tissue

1. Прорастание сосудов. Growth of vessels
2. Формирование рубца из грануляционной соединительной ткани (коллаген I типа)
Formation of scar from granulation connective Tissue (collagen 1 type)

Хрящевая ткань

Cartilage tissue

Клетки (cells):

Хондроциты **Хондробласты**
Chondrocyte ***Chondroblast***

Межклеточное вещество
Intercellular substance:

Основное вещество – **Хондромукоид**
(amorphous substance):


- 80% воды (***water***)
- 4-7 % минеральных солей (***salts***)
- белки – гликопротеиды, протеогликаны, альбумин, ***glycoproteins, proteoglycans, albumin***
- ГАГ (***GAG***) - гиалуроновая кислота, хондроитинсульфаты

**Хондриновые
волокна:**

- коллагеновые
collagenic
- эластические
elastic

Гиалиновая хрящевая ткань

Hyaline cartilage tissue



Волокнистый слой

Fibrous layer

Хондрогенный слой

Chondrogenic layer

Хондробласты

Chondroblasts

Молодые

Хондроциты

Young

chondrocytes

Межтерриториальные

зоны

Interterritorial matrix

Надхрящница

Perichondrium

Зона молодого

хряща

Zone of young

chondrocytes

Зона зрелого

хряща

Zone of mature

chondrocytes

Изогенные группы

Isogenic group

Зрелые хондроциты

mature chondrocytes

Клеточная

Территория

Territorial cell matrix

The cartilage grows in two ways:


- interstitial growth (inside) by formation **isogenic groups**,
- appositional growth (outside) by stratification new ***young chondrocytes***.

Виды хрящевой ткани

Types of cartilage tissue


Гиалиновый (**hyaline**) хрящ

- a lime- CaCO_3 - is put inside,
- collagenic fibres 2 type, thin.




Эластический (**elastic**) хрящ

- a lime is not put inside,
- very much elastic fibres



Волокнистый (**fibrous**)

хрящ (thick bunches of collagenic fibres passing in a sinew or a bone).



Изогенные группы (Isogenous groups)