

Западно-Казахстанский государственный медицинский университет имени
М.Оспанова

Тема: Влияние неинвазивной стимуляции блуждающего нерва на частоте атаки с течением времени, так и расширенные показатели ответа у пациентов с хронической кластерной головной болью

Подготовила Тыныштыкбаева А. У

Актуальность

Кластерная головная боль (СН) значительное время накладывается здравоохранения и затрагивает приблизительно 1 в 1000 человек во всем мире, с хронической кластерной головной боли (ССН) , содержащей от 10% до 15% случаев. Клинические исследования и основанные на фактических данных рекомендации по профилактике СН ограничены, и некоторые профилактические варианты лечения этого заболевания.

- Кластерная головная боль - самый частый вид тригеминальных вегетативных цефалгий, который характеризуется уникальным видом головной боли в сочетании со столь же уникальными сопровождающими симптомами.

Цель

- Профилактика и лечение ХКГЛ
- - изучение неинвазивной стимуляции блуждающего нерва и сравнение его со стандартной терапией, чтобы показать значительные результаты лечения

Вопрос

- Следует ли назначать неинвазивную стимуляцию блуждающего нерва пациентам с хронической кластерной головной болью для снижения частоты атак ?

PICO

- Р-пациенты с хронической кластерной головной болью
- I- Неинвазивная стимуляция блуждающего нерва
- C-стандартное лечение
- O- снижение частоты атак

Дизайн исследования

- многоцентровое, проспективное, рандомизированное, контролируемое исследование

Выборка

- Удобная

Задачи

- ◆ **1. Отобрать 100 пациентов с кластерной головной болью в возрасте от 18 до 70 лет.**
- 2. Разделить на две группы, в каждой из которых по 50 пациентов:**
 - ◆ **- основная группа (неинвазивная стимуляция блуждающего нерва)**
 - ◆ **- контрольная группа (стандартное лечение).**

Критерии включения

- ✓ Пациенты в возрасте от 18 до 70 лет
- ✓ Пациенты страдающие хронической кластерной головной болью

Критерии исключения

- Внутрочерепные, сонные аневризмы, кровоизлияние,
- хирургическое вмешательство (например эндартерэктомией или сосудистой хирургии шеи),
- обмороки или судороги
- травмы головы.
- сердечно - сосудистые заболевания
- ток имплантация с электрическими или нейростимуляционными устройствами или металлоизделиями.

Этические аспекты

- **Одобрено КЭ**
- **Информированное согласие с полным раскрытием всей необходимой информации**
- **Имеют право отказаться от исследования в любое время**
- **Эквивалентность**
- **Действие в интересах пациента**
- **Полезность для пациента и общества**

VAGUS NERVE STIMULATION INHIBITS CORTICAL SPREADING DEPRESSION

The publisher's final edited version of this article is available at [Pain](#)

See other articles in PMC that [cite](#) the published article.

[Go to:](#)

Abstract

Vagus nerve stimulation has recently been reported to improve symptoms of migraine. Cortical spreading depression is the electrophysiological event underlying migraine aura, and a trigger for headache. We tested whether vagus nerve stimulation inhibits cortical spreading depression to explain its anti-migraine effect. Vagus nerve stimulation was delivered either non-invasively through the skin or directly by electrodes placed around the vagus nerve unilaterally. Systemic physiology was monitored throughout the study. Both non-invasive transcutaneous and invasive direct vagus nerve stimulation significantly suppressed spreading depression susceptibility in the occipital cortex in rats. The electrical stimulation threshold to evoke a spreading depression was elevated by more than two-fold, the frequency of spreading depressions during continuous topical 1M KCl was reduced by ~40%, and propagation speed of spreading depression was reduced by ~15%. This effect developed within 30 minutes after vagus nerve stimulation. Non-invasive transcutaneous vagus nerve stimulation was as efficacious as direct invasive vagus nerve stimulation, and the efficacy did not differ between the ipsilateral and contralateral hemispheres. Our findings provide a potential mechanism by which vagus nerve stimulation may be efficacious in migraine, and suggest that susceptibility to spreading depression is a suitable platform to optimize its efficacy.

Keywords: Migraine, Cortical spreading depression, Vagus nerve stimulation

[Go to:](#)

1. Introduction

Migraine is a highly prevalent and disabling disease with tremendous socioeconomic impact. Despite intense research, therapeutic options are limited. Even the most effective acute abortive drugs, triptans, can only achieve a 30% two-hour pain-free response rate at best [28]. Considering the complex pathophysiology and genetics of migraine [2; 52; 54], new therapeutic modalities are needed for patients who do not respond to currently available treatments.

Vagus nerve stimulation (VNS) is an FDA-approved treatment for seizures and depression, currently under investigation as a non-pharmacological therapy in migraine. Patients with refractory epilepsy or major depression have reported remarkable improvement in their migraines after VNS [11; 17; 34; 43; 59], and a small case series suggested efficacy in chronic refractory migraine [48]. In recent open-label, single-arm, multiple-attack studies, VNS afforded a 22% two-hour pain-free rate for moderate or severe attacks [10; 30]. Despite these promising early results, the mechanism of action of VNS in migraine is unknown, hampering the efforts to optimize its efficacy and minimize potential side effects.

Cortical spreading depression (CSD) is a slowly propagating wave of neuronal and glial depolarization that underlies migraine aura. Once triggered, CSD activates downstream inflammatory and nociceptive pathways leading to headache that can last many hours and sometimes days [6; 15; 37; 64; 65]. Animal models of CSD have been widely used as a validated platform for screening pharmacological therapies for migraine [5; 7]. Indeed, migraine prophylactic drugs belonging to different pharmacological classes all inhibit CSD as a common mode of action [7; 23], suggesting that CSD is a headache trigger and a viable therapeutic target in migraine.

Previous work in animal models has shown that VNS activates nucleus tractus solitarius (NTS), locus ceruleus (LC) and dorsal raphe nuclei (DRN) [4; 21; 22; 32], all of which can suppress CSD susceptibility by enhancing serotonergic or β -adrenergic activity [29; 42; 46; 56]. VNS has been effective for the treatment of seizures and depression which show a strong clinical association with migraine [27], and many migraine prophylactic drugs are antiepileptics or antidepressants, and suppress CSD [5]. Moreover, VNS has reproducibly shown efficacy in animal models of ischemic [3; 33; 35] or traumatic brain injury [61], where CSD plays a major pathophysiologic role [40]. We, therefore, hypothesized that VNS suppresses CSD susceptibility as one mechanism of action in migraine, and tested this electrophysiologically in previously validated animal models under full systemic physiological monitoring.

[Go to:](#)

2. Methods

2.1 Ethics statement

Experiments were carried out in accordance with the Guide for Care and Use of Laboratory Animals (National Institutes of Health Publication No. 85-23, 1996), and approved by the institutional review board (Massachusetts General Hospital Subcommittee on Research Animal Care).

Вопрос

Эффективно ли назначение инвазивной стимуляции блуждающего нерва для улучшения симптомов мигрени?

PICO

- ✓ **P-** Больные с мигренью.
- ✓ **I-** инвазивная стимуляция блуждающего нерва
- ✓ **C-** *стандартное лечение*
- ✓ **O-** улучшение симптомов мигрени

Список литературы

1. <https://ru.wikipedia.org/wiki/>.
2. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4943574/>