

*Kazakh National Medical
University
after name Asfendiyarov*



*С.Ж.Асфендияров
атындағы
Қазақ Ұлттық Медицина
Университеті*

Department of foreign languages

Pulmonary Tuberculosis

Faculty: GM

Course: II

Group: 15-027-2

Checked: Saiynova M.B.


Student: Erzhan B.E.

PLAN

- Pulmonary Tuberculosis
- Epidemiology
- Incidence
- Transmission
- Diagnostics
- Chest radiography
- Signs and Symptoms
- Vaccines
- Current Surgical Intervention

Pulmonary Tuberculosis

- **Tuberculosis** (abbreviated as **TB** for *tubercle bacillus* or **Tuberculosis**) is a common and often deadly [infectious disease](#) caused by [mycobacteria](#), mainly [Mycobacterium tuberculosis](#). Tuberculosis usually attacks the lungs (as [pulmonary](#) TB).
- Scanning electron micrograph of Mycobacterium tuberculosis




Epidemiology

- According to the World Health Organization (WHO), nearly 2 billion people—one third of the world's population—have been exposed to the tuberculosis pathogen.
- Annually, 8 million people become ill with tuberculosis, and 2 million people die from the disease worldwide.
- In 2004, around 14.6 million people had active TB disease with 9 million new cases.
- The annual [incidence](#) rate varies from 356 per 100,000 in [Africa](#) to 41 per 100,000 in the [Americas](#).
- Tuberculosis is the world's greatest infectious killer of women of reproductive age and the leading cause of death among people with [HIV/AIDS](#).
- Most common infectious cause of death worldwide
- Latent phase of TB enabled it to spread to one third of the world population
- 8,000,000 new cases each year
- 3,000,000 infected patients die

Epidemiology

Global incidence of TB



Major changes in trends secondary to HIV

- 1953-1985 cases decreased from 84,304 to 22,201

- during this period cases were reactivation of old infection and elderly


- TB and AIDS registries suggests that HIV-infected pts account for 30-50% increase in cases of TB

Incidence

- 1985-1990 TB cases increased 55% in Hispanics and 27% in African Americans
- Populations at risk
 - Foreign-born individuals
 - Low socioeconomic status
 - Cancer pts
 - Celiac disease
 - Cigarette smokers
 - TNF- α antagonists
 - Corticosteroids
- - HIV

Transmission


- When people suffering from active pulmonary TB cough, sneeze, speak, or spit, they expel infectious [aerosol](#) droplets 0.5 to 5 [μm](#) in diameter.
- A single sneeze can release up to 40,000 droplets.
- People with prolonged, frequent, or intense contact are at particularly high risk of becoming infected, with an estimated 22% infection rate.
- A person with active but untreated tuberculosis can infect 10–15 other people per year.
- Others at risk include people in areas where TB is common,



Transmission

- people who inject drugs using unsanitary needles,
- residents and employees of high-risk congregate settings,
- medically under-served and low-income populations,
- high-risk racial or ethnic minority populations,
- children exposed to adults in high-risk categories,
- patients immunocompromised by conditions such as HIV/AIDS, people who take immunosuppressant drugs,
- and health care workers serving these high-risk clients.

Pathogenesis




Diagnostics

- Inject intradermally 0.1 ml of 5TU PPD tuberculin
- Produce wheal 6 mm to 10 mm in diameter
- Represent DTH (delayed type hypersensitivity)



Reading of Mantoux test

- Read reaction 48-72 hours after injection
- Measure only induration
- Record reaction in mm



Classifying the Tuberculin Reaction

>5 mm is classified as positive in

- HIV-positive persons
- Recent contacts of TB case
- Persons with fibrotic changes on CXR consistent with old healed TB
- Patients with organ transplants and other immunosuppressed patients

>10 mm is classified as positive in

Recent arrivals from high-prevalence countries

Injection drug users

Residents and employees of high-risk settings

Mycobacteriology laboratory personnel

Persons with clinical conditions that place them at high risk

Children <4 years, or children and adolescents exposed to adults in high-risk categories

>15 mm is classified as positive in

Persons with no known risk factors for TB

Factors may affect TST

- **False negative**


- Faulty application
- Anergy
- Acute TB (2-10 wks to convert)
- Very young age (< 6 months old)
- Live-virus vaccination
- Overwhelming TB disease

- **False positive**

- BCG vaccination (usually <10mm by adulthood)
- Nontuberculous mycobacteria infection

Chest Radiography

- Abnormalities often seen in apical or posterior segments of upper lobe or superior segments of lower lobe
- May have unusual appearance in HIV-positive persons
- **Cannot confirm diagnosis of TB!!**



Chest radiography

- No chest X-ray pattern is absolutely typical of TB
- 10-15% of culture-positive TB patients not diagnosed by X-ray
- 40% of patients diagnosed as having TB on the basis of x-ray alone do not have active TB

Specimen Collection

- **Obtain 3 sputum specimens for smear examination and culture**
- Persons unable to cough up sputum
 - induce sputum
 - bronchoscopy
 - gastric aspiration
- Follow infection control precautions during specimen collection

Number of sputum samples required

- overall diagnostic yield for sputum examination related to
 - the quantity of sputum (at least 5 mL)
 - the quality of sputum
 - multiple samples obtained at different times to the laboratory for processing
 - 3 samples obtained at least eight hours apart with at least one sample obtained in the early morning

Smear Examination

- Strongly consider TB in patients with smears containing acid-fast bacilli (AFB)
- Results should be available within 24 hours of specimen collection
- Presumptive diagnosis of TB
- Not specific for *M. tuberculosis*

Signs and Symptoms

- Hemoptysis
- Also known as coughing up blood, it is a symptom of bleeding somewhere in the respiratory tract. Frothy and bright red blood may come from the nose, mouth, or throat (upper respiratory tract), the lower respiratory tract, or the lungs. The seriousness of the disorder depends on the cause of the bleeding.



Signs and Symptoms

- Anorexia
- The symptom of poor appetite whatever the cause



Treatment



Prevention

- TB prevention and control takes two parallel approaches.
- In the first, people with TB and their contacts are identified and then treated.
- Identification of infections often involves testing high-risk groups for TB.
- In the second approach, children are vaccinated to protect them from TB.

Vaccines

- Many countries use [Bacillus Calmette-Guérin](#) (BCG) vaccine as part of their TB control programs, especially for infants. According to the W.H.O., this is the most often used vaccine worldwide, with 85% of infants in 172 countries immunized in 1993.
- BCG provides some protection against severe forms of pediatric TB
- unreliable against adult pulmonary TB,
- Currently, there are more cases of TB on the planet than at any other time in history
- urgent need for a newer, more effective vaccine that would prevent all forms of TB—including drug resistant strains—in all age groups and among people with HIV.

Current Surgical Intervention

- Patients with hemoptysis first received Bronchial Artery Embolization because of the recurrent hemoptysis.
- Current indication of Lung Resection for **pulmonary tuberculosis** includes MDR-TB with a poor response to medical therapy, hemoptysis due to bronchiectasis or Aspergillus superinfection, and destroyed lung as previously reported, which are consistent with our indications.
- Surgery remains a crucial adjunct to medical therapy for the treatment of MDR-TB and medical failure lesions.

The used materials:

- www.google.kz
- www.yandex.kz
- www.slideshare.net
- www.medterms.com
- www.nlm.nih.gov



**Thank you for
the attention**

